


CP -violation and $\pi\pi$ -interaction in the radiative decays of K_L and K_S .

S. S. Bulanov

*Institute of Theoretical and Experimental Physics,
B. Cheremyskinskaya ul., 25, 117259 Moscow, Russia.
e-mail: bulanov@heron.itep.ru*



The phases of terms of $K_{L,S} \rightarrow \pi^+\pi^-\gamma$ decay amplitude that arise from the $\pi\pi$ interaction are obtained by using a simple realistic model of $\pi\pi$ interaction via virtual ρ -meson, instead of the ChPT. It is shown that the standard ChPT approach cannot reproduce the contribution of the ρ -meson to the $\pi\pi$ interaction. The interference between the terms of amplitude with different CP -parity appears only when the photon is polarized (linearly or circularly). Instead of measuring the linear polarization, the angular correlation between the $\pi^+\pi^-$ and e^+e^- planes in $K_{S,L} \rightarrow \pi^+\pi^-e^+e^-$ decay can be studied.

1 Introduction

The theoretical and experimental study of the CP -violation in the radiative decays of the K_L and K_S has a long history. In view of future precise measurements of these decays we have recalculated the above effects. Generally the results that will be presented in this talk are in agreement with the previous ones. A few discrepancies are caused by more realistic evaluation of phases caused by the $\pi\pi$ -interaction in the K_L and K_S decays. More accurate calculation of these phases is the prerequisite for extracting the precise values of the CP -violating parameters in the K -meson decays.

The pattern of the CP -violation in the $K_{L,S} \rightarrow \pi^+\pi^-\gamma$ decays was theoretically predicted in the 1960s^{1, 2, 3, 4}. In the 1990s these decays were thoroughly studied using ChPT^{5, 6}. The $K_L \rightarrow \pi^+\pi^-\gamma$ decay attracted special attention, because the contributions of the K_L decay amplitude terms with opposite CP -parity are of comparable magnitude and this makes the CP -violation to be distinctively seen experimentally^{7, 8} with the result $Br(CP = +1) = (1.49 \pm 0.08) \times 10^{-5}$, $Br(CP = -1) = (3.19 \pm 0.16) \times 10^{-5}$. Contrary to this in the case of the K_S decay the CP -violation is difficult to detect due to the fact that the internal bremsstrahlung contribution shades the contribution of the direct emission⁸.

In this talk I present the results of the calculations of the phases of the amplitude terms, connected with the $\pi\pi$ -interaction, using instead of ChPT a simple realistic model of $\pi\pi$ -interaction via virtual ρ -meson⁹. I compare my results on the photon energy dependence of the interference between the internal bremsstrahlung and the electric direct emission in the K_S decay¹¹ with the results of Ref. ⁵ obtained in the framework of ChPT. It is proved that the "interference branching ratio" differs from the one obtained in Ref. ⁵ (see Table 1) and that the standard ChPT approach cannot reproduce the contribution of the ρ . In order to show that the inclusion of the ρ into the analysis of the $K_{S,L}$ decays is important, I address the problem of P -wave $\pi\pi$ scattering. I compare the experimental data with the results obtained in the framework of different models: a) the standard ChPT, b) the ChPT with ρ and c) the simple realistic model.

According to the approach proposed in Ref. ¹⁰ for K_L decay, I examine the $K_S \rightarrow \pi^+\pi^-\gamma$ decay probability with the polarized photon, taking into account various cases of the photon polarization. It is known that an alternative to measuring the linear photon polarization, the angular correlation of $\pi^+\pi^-$ and e^+e^- planes in $K_{S,L} \rightarrow \pi^+\pi^-e^+e^-$ decay can be studied. I present the result of the calculations of the CP -violating asymmetry in the case of the K_L decay in order to compare our results with those of Ref. ¹⁵. I also present the result for the CP -violating asymmetry in the case of the K_S decay.

2 The amplitude structure

The amplitudes of $K_{S,L}(r) \rightarrow \pi^+(p)\pi^-(q)\gamma(k,e)$ decays are made up of two terms: the internal bremsstrahlung (B) and direct emission (D). In its turn, D is a sum of an electric term(E_D) and a magnetic term(M_D). In accordance with the above, the amplitudes of the $K_{S,L}$ decays can be written as follows

$$A(K_S \rightarrow \pi^+\pi^-\gamma) = eAe^{i\delta_0^0}T_B + e(ae^{\delta_1^1} + be^{\delta_b})T_E + ie\eta_{+-}ce^{\delta_1^1}T_M, \quad (1)$$

$$A(K_L \rightarrow \pi^+\pi^-\gamma) = \eta_{+-}eAe^{i\delta_0^0}T_B + e\eta_{+-}(ae^{\delta_1^1} + be^{\delta_b})T_E + i ece^{\delta_1^1}T_M, \quad (2)$$

where $T_B = \frac{pe}{pk} - \frac{qe}{qk}$, $T_E = (pe)(qk) - (qe)(pk)$, $T_M = \varepsilon_{\mu\nu\rho\sigma}p_\mu q_\nu k_\rho e_\sigma$; δ_0^0 and δ_1^1 are the S -wave and P -wave pion scattering phases respectively and η_{+-} is the well known CP -violation parameter in the $K_L \rightarrow \pi^+\pi^-$ decay. The factor $A \equiv A(K \rightarrow \pi^+\pi^-)$ is determined by the Low theorem for bremsstrahlung¹⁶. As follows from Eqs. (1, 2) E_D consists of two terms. The first term ($ae^{\delta_1^1}$) describes the loops of heavy particles. The second term (be^{δ_b}) describes the loops of pions. Such subdivision is convenient because the pion loops contribution has an absorptive part and hence a phase, contrary to the contribution of the heavy particle loops.

The phases of the pion loops and bremsstrahlung contributions in equations (1), (2) are defined by the strong interaction of pions via virtual ρ -meson:

$$L = -\frac{g}{\sqrt{2}}\varepsilon_{ijk}(\phi^i\partial_\mu\phi^j - \partial_\mu\phi^i\phi^j)B_\mu^k, \quad (3)$$

where i, j, k are isotopic indices, ϕ is the pion field, B_μ^k – ρ -meson field.

In the case of internal bremsstrahlung contribution to the $K_{S,L}$ decay probability the interaction of pions can be described by the diagrams shown in Fig. 1a. Though pions are in P -wave in the final state this group of diagrams results in the δ_0^0 phase of the amplitude. It is due to the fact that the interaction of pions occurs not in the final state, but in the intermediate one.

The emission of the photon from the loops of pions is governed by another group of diagrams shown in Fig. 1b. Each of these diagrams is divergent but their sum is finite. The matrix element arising from the diagrams, shown in Fig. 1b, takes the form:

$$E_D^{loop} = \frac{eg^2A}{\pi^2}F(s)D(s)\frac{T_E}{rk} = ebT_E, \quad (4)$$




Figure 1: a) The interaction of pions in the case of the internal bremsstrahlung. b) The emission of the photon from the loops of virtual particles (D).

with

$$F(s) = \frac{1}{2} + \frac{s}{2rk} [\beta K(\beta) - \beta_0 K(\beta_0)] - \frac{m_\pi^2}{rk} (K^2(\beta) - K^2(\beta_0)) + \frac{i\pi}{2rk} \left(\frac{s}{2} (\beta - \beta_0) - 2m_\pi^2 (K(\beta) - K(\beta_0)) \right), \quad (5)$$

where $s = (r - k)^2$, $\beta = \sqrt{1 - 4m_\pi^2/s}$, $\beta_0 = \sqrt{1 - 4m_\pi^2/m_K^2}$, $K(\beta) = \text{Arth}\left(\frac{1}{\beta}\right)$, $D(s)$ is the ρ propagator, g is the interaction constant of $\rho\pi\pi$.

Note that since $F(s)$ is complex, the phase of the loop contribution is not equal to the pion P -wave scattering phase. Heavy particles in the loop can also contribute to the electric direct emission amplitude, though they do not produce any additional phase. The possible intermediate states are πK , $K\eta$ and KK . However, the KK loop vanishes in the limit $m_{K^0} = m_{K^+}$. The contribution of these loops is calculated using the assumption that the pseudoscalar mesons (π , K , η) couple to ρ with equal strength, i. e. $g_{\rho K\pi} = g_{\rho K\eta} = g_{\rho KK} = g$.

3 Comparison with ChPT

The electric direct emission manifests itself in the decay probability mainly through the interference with the internal bremsstrahlung. Let us compare the results on the "interference branching ratio" in the K_S decay obtained in Ref. ⁵ with the results obtained in the framework of the simple realistic model ¹¹. In the framework of the ChPT the electric direct emission is a sum of the loop and counterterm contributions.

Table 1

cut in ω	$\omega > 20$ Mev	$\omega > 50$ Mev	$\omega > 100$ Mev
10^6 Interf^{11}	-6.3	-4.8	-1.7
$10^6 \text{ Interf}(k_f = 0)^5$	-6.2	-5.0	-2.0
$10^6 \text{ Interf}(k_f = 0.5)^5$	-10.5	-8.3	-3.3
$10^6 \text{ Interf}(k_f = 1.0)^5$	-14.8	-11.7	-4.7
$10^6 \text{ Interf}(k_f = -0.5)^5$	-1.9	-1.6	-0.6
$10^6 \text{ Interf}(k_f = -1.0)^5$	+2.4	+1.8	+0.7

In Table 1 I present interference contributions to the branching ratio of the $K_S \rightarrow \pi^+\pi^-\gamma$ decay, for different values of the ω cut, along with the results of Ref. ⁵. In the case of $k_f = 0$ the "interference branching ratio" obtained here and the one obtained in the framework of the ChPT are in agreement for the photon energy cut $\omega > 20$ MeV. However, for the photon energy cuts $\omega > 50$ MeV and $\omega > 100$ MeV the discrepancy appears. It is due to the fact that the photon spectra of these results differ. It can be clearly seen from Fig. 2a, where my result (solid curve) along with the results of Ref. ⁵ (dashed curves, $k_f = -0.5, 0, +0.5$) for the photon spectra of the interference contribution are shown.

If the counterterm contributions are switched on, then the arising discrepancy is rather large for given in Ref. ⁵ values of the counterterm contributions. It should be stressed that the

counterterm contributions don't depend on the photon energy and exhibit the behavior different from our results (Fig. 2a). The discrepancy mentioned above is due to the fact that in Ref. ⁵ the ρ contribution shows up only in low-energy constants, while the resonance contribution was not considered. The phase of the electric direct emission amplitude was taken to be $\delta_1^1(m_K)$, the phase of P -wave $\pi\pi$ scattering at fixed energy $\sqrt{s} = m_K$. Instead I used the simple realistic model of $\pi\pi$ scattering via ρ -meson taking into account the energy dependence of the $\delta_1^1(s)$ phase. As it can be seen from Fig. 2a the standard ChPT approach even with higher order counterterms taken into account cannot reproduce the contribution of the ρ -meson.




Figure 2: a) The photon energy dependence of the interference contribution. The result of the present paper is represented by solid curve. The results of ChPT – by dashed curves. The upper dashed curve corresponds to $k_f = -0.5$. The lower dashed curve – to $k_f = 0.5$. The dashed curve in between – to $k_f = 0$. b) The energy dependence of the $\pi\pi$ scattering phase $\delta_1^1(s)$ in the P -wave, in degrees. The experimental data are represented by stars. The ChPT without ρ contribution – by diamonds. The ChPT with the ρ contribution taken into account – by filled triangles. The ChPT with ρ in two-loop approximation – by empty triangles. My results – by boxes.

Actually, in the case of $\pi\pi$ scattering in the framework of the ChPT the ρ -meson showed up in two ways: as low-energy constants and as a direct resonance. However, in papers on the $K \rightarrow \pi\pi\gamma$ decays only low-energy constants are accounted for. Such approach doesn't take into account the energy dependence of the $\delta_1^1(s)$ phase, i. e. the behavior of the ρ propagator, because the contribution of low-energy constants doesn't depend on the photon energy. Thus, some dynamical features are missing in the ChPT approach for the $K \rightarrow \pi\pi\gamma$ decays.

In order to show that the inclusion of the ρ into the analysis of the electric direct emission in the $K \rightarrow \pi\pi\gamma$ decays is important, I address the the problem P -wave of $\pi\pi$ scattering. In Fig. 2b I present the behaviour of the P -wave $\pi\pi$ -scattering phase δ_1^1 calculated in the framework of different models: a) the standard ChPT ¹⁷, b) the ChPT with ρ ¹⁸ and c) the simple realistic model. As it is seen from Fig. 2b the result of the standard ChPT is in strict disagreement with the experimental data. It is due to the fact that the ρ contribution was not taken into account. The simple realistic model and the ChPT with ρ are in accordance with the experimental data. Thus, the inclusion of the ρ in the analysis of the $K \rightarrow \pi\pi\gamma$ decays is important.

4 Analysis in terms of Stokes parameters

It is worth mentioning that there is no interference between the amplitude terms with opposite CP -parity, if photon polarization is not observed. However, the interference is nonzero when the polarization is measured. Therefore, any CP -violation involving interference of electric and magnetic amplitudes is encoded in the polarization state of the photon.

In order to study this interference we write the $K_{L,S}$ decay amplitude more generally as

$$A(K_{S,L} \rightarrow \pi^+\pi^-\gamma) = ET_E + MT_M, \quad (6)$$

where for the K_S decay E and M have the form $E = eb + eAe^{i\delta_0^0}/(pk \ qk)$ and $M = ie\eta_{+-}c$, and in the case of the K_L decay we have $E = \eta_{+-} [eb + eAe^{i\delta_0^0}/(pk \ qk)]$ and $M = iec$.

The photon polarization can be defined in the terms of the density matrix

$$\rho = \begin{pmatrix} |E|^2 & E^* M \\ EM^* & |M|^2 \end{pmatrix} = \frac{1}{2}(|E|^2 + |M|^2) [1 + \mathbf{S}\boldsymbol{\tau}], \quad (7)$$

where $\boldsymbol{\tau} = (\tau_1, \tau_2, \tau_3)$ are Pauli matrices, $\mathbf{S}$ is the Stokes vector of the photon with components

$$S_1 = \frac{2\text{Re}(E^* M)}{(|E|^2 + |M|^2)}, \quad S_2 = \frac{2\text{Im}(E^* M)}{(|E|^2 + |M|^2)}, \quad S_3 = \frac{(|E|^2 - |M|^2)}{(|E|^2 + |M|^2)}. \quad (8)$$

In order to obtain a quantitative estimate of the CP -violation effects the photon energy dependence of the Stokes vector components can be studied. In Fig. 3a and 3b the photon energy dependence of the S_1 (coefficient of an interference term in case of linear polarization) and S_2 (net circular polarization) in the $K_{L,S}$ decay respectively are shown. In figure 3c I present the S_3 photon energy dependence in $K_{L,S}$ decays to obtain the estimate of the relative strength of the CP -violation effects in the decays under consideration. The obtained results on the K_L decay coincide with the results of Ref. ¹⁰. Taking into account the physical meaning of the S_3 I conclude that the CP -violation effects in K_S decay are substantially smaller. The reason is that the bremsstrahlung contribution shades the magnetic direct emission contribution even for the high photon energies.




Figure 3: a) Stokes parameters S_1 (upper curve) and S_2 (lower curve) for the K_L decay. b) Stokes parameters S_1 (upper curve) and S_2 (lower curve) for the K_S decay. c) Stokes parameter S_3 for the K_S decay (straight line $S_3 = 1$) and for the K_L decay (lower curve).

It was suggested in Refs. ^{12, 13} to use in place of $\mathbf{e}$ ($e_\mu = (0, \mathbf{e})$), orthogonal to $\mathbf{k}$ ($k_\mu = (\omega, \mathbf{k})$), the vector $\mathbf{n}_1$ normal to the e^+e^- plane in the decay $K_L \rightarrow \pi^+\pi^-e^+e^-$. This can be achieved by replacing e_μ in the radiative amplitude (1, 2) by $e/k^2 \bar{u}(k_-)\gamma_\mu v(k_+)$. This motivates the study of the distribution $d\Gamma/d\phi$ in the decays $K_{S,L} \rightarrow \pi^+\pi^-e^+e^-$, where ϕ is an angle between $\pi^+\pi^-$ and e^+e^- planes.

The resulting CP -violating asymmetry in the K_L decay is $|A_{\pi\pi, ee}^L| = (13.4 \pm 0.9)\%$. It coincides within the accuracy of the calculation with the theoretical prediction $|A_{\pi\pi, ee}^L| = 14\%$, obtained in Refs. ^{10, 12, 13, 15} and the experimental result $(13.6 \pm 2.5 \pm 1.5)\%$ ¹⁴.

The CP -violating asymmetry in the K_S decay is $|A_{\pi\pi, ee}^S| = (5.1 \pm 0.4) \times 10^{-5}$. This value of the asymmetry could be expected qualitatively from the analysis of the $K_S \rightarrow \pi^+\pi^-\gamma$ decay amplitude, where the CP -violating magnetic direct emission contribution is substantially small compared to the CP -conserving part of the amplitude.

5 Conclusions

In the case of the radiative K -meson decays the phases of amplitude terms using a simple realistic model of pion-pion interaction ⁹ were calculated. Also the pion loop contribution (E_D^{loop}) to the electric direct emission amplitude was calculated.

I compared my results on the interference contribution to the $K_S \rightarrow \pi^+\pi^-\gamma$ decay probability with those of Ref. ⁵ and found that the "interference branching ratio" differs from the the one

obtained in Ref. ⁵ (see Table 1). This discrepancy arises from different models of $\pi\pi$ interaction and the fact that I took into account the energy dependence of the phases, while in the standard ChPT approach the phases were taken at the fixed energy $\sqrt{s} = m_K$. Thus it is clear that the standard ChPT approach for the $K \rightarrow \pi\pi\gamma$ decays even with higher order counterterms taken into account cannot reproduce the contribution of the ρ meson.

In order to show that the inclusion of the ρ meson into the analysis of the electric direct emission in the $K \rightarrow \pi\pi\gamma$ decays is important, I addressed the problem of the $\pi\pi$ scattering in P -wave. I compared with the experimental data the predictions for the phase of the $\pi\pi$ scattering in the P -wave obtained in the framework of different models. As seen from Fig. 2b the simple realistic model and the ChPT with ρ are in accordance with the experimental data. The ChPT without ρ shows strong disagreement with data. Thus the inclusion of the ρ meson, done in the framework of the simple realistic model, in the $K \rightarrow \pi\pi\gamma$ decays is important.

Regarding the dependence of the K_S decay probability on photon polarization I found that the effects of CP -violation are small compared to that of the K_L decay, which could be qualitatively expected from the analysis of the K_S decay. The reason is that the bremsstrahlung contribution shades the magnetic direct emission contribution even for the high photon energies.

I also studied the $K_{S,L} \rightarrow \pi^+\pi^-e^+e^-$ decays. The central values of the asymmetries presented in this talk, obtained in Refs. ^{10, 12, 13, 15} and measured experimentally ¹⁴ coincide within the accuracy of the calculation. I found that the CP -violating asymmetry in the case of the K_S decay is substantially smaller than in the K_L case, as it could be expected from the analysis of the $K_S \rightarrow \pi^+\pi^-\gamma$ decay.

The author appreciates L. B. Okun's scientific supervision and formulation of this problem. The author would also like to thank R. B. Nevzorov for fruitful discussions and G. D'Ambrosio, M. I. Vysotsky, E. P. Shabalin for valuable remarks and the Organizing Committee for the warm and stimulating atmosphere of the Moriond Conference.

References

1. H. Chew, Nuovo Cimento **26** 1109 (1962).
2. G. Costa, P. K. Kabir, Nuovo Cimento **A51** 564 (1967).
3. L. M. Sehgal, L. Wolfenstein, Phys. Rev. **162** 1362 (1967).
4. A. D. Dolgov, L. A. Ponomarev, Yad Fiz. **4** 262 (1967).
5. G. D'Ambrosio and G. Isidori, Z. Phys. **C65** 649 (1995).
6. J. Tandean, G. Valencia, submitted to Phys. Rev. D, hep-ph/0008238.
7. A. S. Carrol *et al.*, Phys. Rev. Lett. **44** 529 (1980).
8. E. J. Ramberg *et al.*, Phys. Rev. Lett. **70** 2525 (1993).
9. B. W. Lee, M. T. Vaughn, Phys. Rev. Lett. **4** 578 (1960).
10. L. M. Sehgal, J. van Leusen, Phys. Rev. Lett. **83** 4933 (1999).
11. S. S. Bulanov, submitted to Nucl. Phys. B, hep-ph/0109025.
12. L. M. Sehgal, M. Wanninger Phys. Rev. D**46** 1035 (1992).
13. P. Heiliger, L. M. Sehgal, Phys. Rev. D**48** 4146 (1993).
14. A. Alavi-Harati *et al.*, Phys. Rev. Lett. **84** 408 (2000), hep-ex/9908020.
15. J. K. Elwood, M. B. Wise, M. J. Savage, J. W. Walden, Phys. Rev. D**53** 4078 (1996).
16. F. E. Low, Phys. Rev. **110** 974 (1958).
17. J. Gasser, H. Leutwyler, Ann. of Phys. **158** 142 (1984).
18. J. Bijnens *et al.*, Phys. Lett. B**374** 210 (1996).